# How to test for Xylazine

#### **Powders & Pills guide**

### Step 1: Prepare Sample

#### **Crush Your Pill**

 Make sure your pill or powder is crushed finely into a uniform consistency



### Step 2: Choose a Testing Method

# **Test Everything**

- Pour powder into a cup
- Use 1/2 cup or 100ml of water per gram of material
- Mix until fully dissolved

# **Test Bag Residue**

- Make sure the contents inside the bag are a uniform powder
- Dump the contents into another container
- Fill the empty bag with water and mix if necessary

# **Test a Scoop**

- Put one micro scoop of substance into a small cup
- Mix 1tsp or 5ml of water for each scoop of substance

# Option 1 (Most Accurate)



#### **Option 2**




#### **Option 3**



# Step 3: Testing

### **Use the Test Strip**


- Place the test strip wavy side down in the water for 15s. Make sure the water is up to the line
- Remove strip from water and place it on a flat surface for 1-2 minutes




### Step 4: Results

#### **Read the Results**

 A test POSITIVE for xylazine is indicated by ONE RED LINE



- A test NEGATIVE for xylazine is indicated by TWO RED LINES
- Remember, no test is 100% accurate,
  & a negative test does not necessarily
  mean you are safe



NOTE: It should be considered a NEGATIVE test even if a second red line is faded or faint If possible, use an additional test to confirm your results



# How to test for Xylazine

#### **IV Users Guide**

#### Lower Your Risk of Overdose

#### **Be Careful**

- Go slow by taking smaller amounts
- Use with people you trust and inject at different times
- Have Naloxone available and know how to use it



### Step 1: Testing Cooker Residue

### **Prepare Shot**

- Prepare your shot as normal and set aside the needle to test your substance left over in the cooker before injection
- Add enough water to fill your cooker.
  Stir if necessary



# Step 2: Testing

#### **Use the Test Strip**


- Place the test strip wavy side down in the water for 15s. Make sure the water is up to the line
- Remove strip from water and place
  it on a flat surface for 1-2 minutes




# Step 3: Results

#### **Read the Results**

- A test POSITIVE for xylazine is indicated by ONE RED LINE
- A test NEGATIVE for xylazine is indicated by TWO RED LINES



Remember, no test is 100% accurate,
 & a negative test does not necessarily mean you are safe



NOTE: It should be considered a NEGATIVE test even if a second red line is faded or faint If possible, use an additional test to confirm your results

